

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID: SSSPTA1623PAZ

PASSWORD :

TERMINAL (ENTER 1, 2, 3, OR ?):2

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer Agreement.

agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

FILE 'HOME' ENTERED AT 06:25:36 ON 19 JUL 2005

=> file reg
COST IN U.S. DOLLARS . SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST . 0.84 . 0.84

FILE 'REGISTRY' ENTERED AT 06:27:51 ON 19 JUL 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 17 JUL 2005 HIGHEST RN 855596-49-5
DICTIONARY FILE UPDATES: 17 JUL 2005 HIGHEST RN 855596-49-5

New CAS Information Use Policies, enter **HELP USAGE TERMS** for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005


Please note that search-term pricing does apply when conducting SmartSELECT searches.

*
* The CA roles and document type information have been removed from *
* the IDE default display format and the ED field has been added, *
* effective March 20, 2005. A new display format, IDERL, is now *
* available and contains the CA role and document type information. *
*

Structure search iteration limits have been increased. See **HELP SLIMITS** for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>

=> Uploading C:\Documents and Settings\PZucker\My Documents\Examination Auxillary files\10510024\10510024 unsubstituted core.str



```

chain nodes :
7 8 9 10 11 12 13 14
ring nodes :
1 2 3 4 5 6
chain bonds :
6-7 7-8 8-9 9-10 10-11 11-12 12-13 12-14
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6
exact/norm bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 8-9 10-11 11-12 12-14
exact bonds :
6-7 9-10 12-13

```

```

Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:CLASS 12:CLASS 13:CLASS 14:CLASS


```

L1 STRUCTURE UPLOADED

```

=> d 11
L1 HAS NO ANSWERS
L1           STR

```



Structure attributes must be viewed using STN Express query preparation.

```

=> search 11 sss sam
SAMPLE SEARCH INITIATED 06:28:12 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 14803 TO ITERATE

```

```

13.5% PROCESSED       2000 ITERATIONS                           50 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

```

```

FULL FILE PROJECTIONS:   ONLINE   **COMPLETE**
                          BATCH    **COMPLETE**
PROJECTED ITERATIONS:    288773 TO   303347
PROJECTED ANSWERS:       8443 TO    11095

```

L2 50 SEA SSS SAM L1

```

=> d scan

```


```

L2   50 ANSWERS   REGISTRY COPYRIGHT 2005 ACS on STN
IN   2-Propenoic acid, (1S,2E)-1-[(benzoyloxy)methyl]-4-bromo-2-butenoyl ester

```

(9CI)
MF C15 H15 Br O4

Absolute stereochemistry. Rotation (+).
Double bond geometry as shown.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> d cost COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
CONNECT CHARGES	0.74	1.34
NETWORK CHARGES	0.12	0.36
-----		-----
FULL ESTIMATED COST	0.86	1.70

IN FILE 'REGISTRY' AT 06:28:49 ON 19 JUL 2005

=>
Uploading C:\Documents and Settings\PZucker\My Documents\Examination Auxillary
files\10510024\10510024 unsubstituted core2.str




chain nodes :
7 8 9 10 11 12 13 14
ring nodes :
1 2 3 4 5 6
chain bonds :
6-7 7-8 8-9 9-10 10-11 11-12 12-13 12-14
ring bonds :
1-2 1-6 2-3 3-4 4-5 5-6
exact/norm bonds :
1-2 1-6 2-3 3-4 4-5 5-6 7-8 8-9 10-11 11-12 12-14
exact bonds :
6-7 9-10 12-13

Hydrogen count :
1:>= minimum 2 2:>= minimum 2 3:>= minimum 2 5:>= minimum 2 6:>= minimum 1
Match level :
1:Atom 2:Atom 3:Atom 4:Atom 5:Atom 6:Atom 7:CLASS 8:CLASS 9:CLASS 10:CLASS
11:CLASS 12:CLASS 13:CLASS 14:CLASS

L3 STRUCTURE UPLOADED

=> d 13
L3 HAS NO ANSWERS
L3 STR



Structure attributes must be viewed using STN Express query preparation.

=> search l3 sss sam
SAMPLE SEARCH INITIATED 06:30:36 FILE 'REGISTRY'
SAMPLE SCREEN SEARCH COMPLETED - 14803 TO ITERATE

13.5% PROCESSED 2000 ITERATIONS 0 ANSWERS
INCOMPLETE SEARCH (SYSTEM LIMIT EXCEEDED)
SEARCH TIME: 00.00.01

FULL FILE PROJECTIONS: ONLINE **COMPLETE**
BATCH **COMPLETE**
PROJECTED ITERATIONS: 288773 TO 303347
PROJECTED ANSWERS: 0 TO 0

L4 0 SEA SSS SAM L3

=> search l3 sss full
FULL SEARCH INITIATED 06:31:36 FILE 'REGISTRY'
FULL SCREEN SEARCH COMPLETED - 296742 TO ITERATE


100.0% PROCESSED 296742 ITERATIONS 74 ANSWERS
SEARCH TIME: 00.00.02

L5 74 SEA SSS FUL L3

=> d scan

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 2-Butenoic acid, 2-[(1S)-1-[(1R)-3,3-dimethylcyclohexyl]ethoxy]-2-
methylpropyl ester, (2E)- (9CI)
MF C18 H32 O3


Absolute stereochemistry.
Double bond geometry as shown.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 1-Propanol, 2-methyl-2-[1-(3-methylcyclohexyl)ethoxy]-, acetate (9CI)
MF C15 H28 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Propanoic acid, 2,3,3,3-tetrafluoro-2-(heptafluoropropoxy)-, 2-(cyclohexylmethoxy)propyl ester (9CI)
MF C16 H19 F11 O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Acetic acid, (1-oxopropoxy)-, (1R)-1-[(1S)-3,3-dimethylcyclohexyl]ethyl ester (9CI)
MF C15 H26 O4


Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 1-Propanol, 2-[1-(3,3-dimethylcyclohexyl)ethoxy]-2-methyl-, propanoate,
(R*,S*)- (9CI)
MF C17 H32 O3


Relative stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 1,3-Cyclohexanedicarboxylic acid, mono[2-[(2-methyl-1-oxo-2-
propenyl)oxyethyl] ester, homopolymer (9CI)
MF (C14 H20 O6)x
CI PMS

CM 1




L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Cyclohexanecarboxylic acid, 2-hydroxyethyl ester, acetate (8CI)
MF C11 H18 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 2-Propanol, 1-(1-cyclohexyl-1-methylethoxy)-, acetate (9CI)
 MF C14 H26 O3




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN.
 IN 1-Propanol, 2-(1-cyclohexylpropoxy)-2-methyl-, propanoate (9CI)
 MF C16 H30 O3




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 2-Butenoic acid, 2-[1-(3,3-dimethylcyclohexyl)ethoxy]-2-oxoethyl ester
 (9CI)
 MF C16 H26 O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Acetic acid, (1-oxoproxy)-, 1-(3,3-dimethylcyclohexyl)ethyl ester (9CI)
 MF C15 H26 O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 2-Propanol, 1-[1-(3,3-dimethylcyclohexyl)ethoxy]-2-methyl-, propanoate
 (9CI)
 MF C17 H32 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Decanoic acid, 2-hydroxyethyl ester, cyclohexanecarboxylate (8CI)
 MF C19 H34 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 2-Propenoic acid, 2-hydroxyethyl ester, polymer with α -[5-
 [(dimethylamino)thioxomethyl]thio]-1-oxopentyl]- ω -methoxypoly(oxy-
 1,2-ethanediylloxycarbonyl-1,3-cyclohexanediylcarbonyl) and
 octahydro-4,7-methano-1H-inden-5-yl 2-propenoate, diblock (9CI)
 MF (C13 H18 O2 . (C10 H14 O4)n C9 H17 N O2 S2 . C5 H8 O3)x
 CI PMS


CM 1




CM 2



CM 3




L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Butanoic acid, 2-[1-(3,3-dimethylcyclohexyl)-1-methylethoxy]-2-oxoethyl
ester (9CI)
MF C17 H30 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 1-Propanol, 2-(1-cyclohexylethoxy)-2-methyl-, acetate (9CI)
 MF C14 H26 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Butanoic acid, 2-[1-(3,3-dimethylcyclohexyl)ethoxy]-2-oxoethyl ester (9CI)
 MF C16 H28 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 1H-Pyrazole-3-carboxylic acid, 4-[[4-[[2-[2-(cyclohexylmethoxy)ethoxy]1-methyl-2-oxoethoxy]carbonyl]phenyl]azo]-4,5-dihydro-5-oxo-1-phenyl-, 1-methylbutyl ester (9CI)
 MF C34 H42 N4 O8

PAGE 1-A




PAGE 2-A




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 2-Propanol, 1-[1-(3,3-dimethylcyclohexyl)ethoxy]-, acetate (9CI)
MF C15 H28 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Heptanoic acid, 2-hydroxyethyl ester cyclohexanecarboxylate (8CI)
 MF C16 H28 O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN 2-Propenoic acid, 2-[1-[(1R)-3,3-dimethylcyclohexyl]ethoxy]-2-methylpropyl
 ester (9CI)
 MF C17 H30 O3


Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):10


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Acetic acid, (1-oxopropoxy)-, 1-(3,3-dimethylcyclohexyl)-1-methylethyl
 ester (9CI)
 MF C16 H28 O4



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN D-glycero-D-galacto-Non-2-enonic acid, 5-(acetylamino)-2,6-anhydro-4-
 [[bis[[1,1-dimethylethoxy]carbonyl]amino]methylene]amino]-7-O-
 (cyclohexylmethyl)-3,4,5-trideoxy-, methyl ester, 8,9-diacetate (9CI)
 MF C34 H54 N4 O13


Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
 IN Ethanedioic acid, 2-[(1S)-1-[(1R)-3,3-dimethylcyclohexyl]ethoxy]-2-
 oxoethyl methyl ester (9CI)
 MF C15 H24 O6


Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Cyclohexanecarboxylic acid, 2-(acetyloxy)-3-phenoxypropyl ester, (S)-
(9CI)
MF C18 H24 O5

Absolute stereochemistry.




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 1-Propanol, 2-[1-(3,3-dimethylcyclohexyl)ethoxy]-, acetate (9CI)
MF C15 H28 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Cyclohexanecarboxylic acid, 2-hydroxyethyl ester butyrate (8CI)
MF C13 H22 O4




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 2-Propenoic acid, 2-[(1S)-1-[(1R)-3,3-dimethylcyclohexyl]ethoxy]-2-
methylpropyl ester (9CI)


MF C17 H30 O3

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT


L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN 1-Propanol, 2-(1-cyclohexyl-1-methylethoxy)-, propanoate (9CI)
MF C15 H28 O3



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Cyclohexanecarboxylic acid, (1R)-1-[(acetoxy)methyl]-2-(phenylmethoxy)ethyl ester (9CI)
MF C19 H26 O5


Absolute stereochemistry. Rotation (-).



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

L5 74 ANSWERS REGISTRY COPYRIGHT 2005 ACS on STN
IN Acetic acid, (1-oxoproxy)-, (1S)-1-[(1R)-3,3-dimethylcyclohexyl]ethyl ester (9CI)
MF C15 H26 O4

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

HOW MANY MORE ANSWERS DO YOU WISH TO SCAN? (1):0

=> save temp 15 rawfinds/a
ANSWER SET L5 HAS BEEN SAVED AS 'RAWFINDS/A'

=> file caplus
'CAPLUS' IS NOT A VALID FILE NAME
SESSION CONTINUES IN FILE 'REGISTRY'
Enter "HELP FILE NAMES" at an arrow prompt (=>) for a list of files
that are available. If you have requested multiple files, you can
specify a corrected file name or you can enter "IGNORE" to continue
accessing the remaining file names entered.

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	165.20	166.04

FILE 'CAPLUS' ENTERED AT 06:33:54 ON 19 JUL 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2005 VOL 143 ISS 4
FILE LAST UPDATED: 18 Jul 2005 (20050718/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> 15
L6 21 L5

=> save temp odorrefs/a

ENTER L#, L# RANGE, ALL, OR (END):16
ANSWER SET L6 HAS BEEN SAVED AS 'ODORREFS/A'

=> odor

64767 ODOR
11508 ODORS
L7 69532 ODOR
(ODOR OR ODORS)

=> 16 and 17

L8 7 L6 AND L7

=> d 18 1-7 ti

L8 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of alicyclic ester with musk fragrance for perfumes or body care products

L8 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Odorous compositions containing (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methyl-1-propanoate esters

L8 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Methods for the production of novel alicyclic esters having a musky smell

L8 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of cycloalkanecarboxylic acid derivatives for use as fragrances with musk characteristics

L8 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI A method for the calculation of odor character from molecular structure

L8 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of esters with musky odor and their use in perfumery

L8 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Synthesis of ethylene glycol mixed esters of alkanoic and cyclohexanoic acids

=> d 18 1-7 ti fbib abs

L8 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of alicyclic ester with musk fragrance for perfumes or body care products
AN 2005:120865 CAPLUS
DN 142:198227
TI Preparation of alicyclic ester with musk fragrance for perfumes or body care products
IN Eh, Marcus
PA Symrise GmbH & Co. KG, Germany
SO PCT Int. Appl., 42 pp.
CODEN: PIXXD2
DT Patent
LA German
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2005012222	A1	20050210	WO 2004-EP51361	20040705
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY,				

TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW
 RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM,
 AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,
 EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,
 SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,
 SN, TD, TG

DE 2003-10335053 A 20030731

DE 10335053


A1 20050224

DE 2003-10335053


20030731

OS CASREACT 142:198227; MARPAT 142:198227


GI




I



II



III



IV

AB A compds. I [R1 = (un)substituted, (un)branched C1-5-alkyl, (un)branched C2-5-alkylene, C3-5-cycloalkyl] and II [dotted line may be either a single or a double bond] are disclosed. The procedure for the preparation of I comprises the esterification with alcs. III and IV. Thus, propionic acid 2-methyl-2-[1-(2-methylcyclohexyl)ethoxy]propyl ester [II; R1 = Et, dashed line = single bond] was prepared from MeCOCH₂CH₂Me-2 via hydrogenation with Ru/C, reaction with isobutylene oxidation in the presence of BF₃·OEt₂, and acylation with (EtCO)₂O in the presence of Et₃N and catalytic DMAP. A perfume formulation containing I (R1 = Et, dashed line = single bond) was prepared

RE.CNT 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
 TI Odorous compositions containing (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methyl-1-propanol alkenoate esters
 AN 2004:780341 CAPLUS
 DN 141:301056
 TI Odorous compositions containing (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methyl-1-propanol alkenoate esters
 IN Williams, Alvin Scott
 PA Switz.
 SO U.S. Pat. Appl. Publ., 6 pp.
 CODEN: USXXCO.

DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	US 2004186043	A1	20040923	US 2004-792375 WO 2003-IB1079	20040302 A 20030319
	EP 1459735	A1	20040922	EP 2004-100807 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK	20040301
	JP 2004285357	A2	20041014	JP 2004-81377 WO 2003-IB1079	20040319 A 20030319

OS MARPAT 141:301056

AB The present invention relates to the field of perfumery, and more precisely to a compound such as (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methylpropyl 2-propenoate (I). The present invention concerns also the use of the compound in the perfumery industry as well as the compns. or articles associated with the compound. Thus, to a suspension of NaH (0.368 mol) in 1000 mL of dry THF at 0° were added dropwise 70.0 g (1S,1'R)-2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methyl-1-propanol followed by the addition of acryloyl chloride. A 96% pure 2-[1-(3',3'-dimethyl-1'-cyclohexyl)ethoxy]-2-methylpropyl 2-propenoate in the form of a mixture containing 81% of the (1S,1'R) diastereoisomer and 19% of 3 other diastereomers was obtained.. The addition of 60 parts by weight of I to a standard perfuming composition imparted to the latter a powerful musky aspect, which is also smooth and fresh, together with a green, pear's peel connotation and a Galbanum undernote.

L8 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

TI Methods for the production of novel alicyclic esters having a musky smell
 AN 2003:796643 CAPLUS

DN 139:307907

TI Methods for the production of novel alicyclic esters having a musky smell

IN Eh, Marcus

PA Symrise GmbH & Co. KG, Germany

SO PCT Int. Appl., 45 pp.

CODEN: PIXXD2


DT Patent

LA German


FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003082799	A2	20031009	WO 2003-EP3294	20030329
	WO 2003082799	A3	20031231		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	DE 10214675	A1	20031016	DE 2002-10214675	A 20020403
	BR 2003004218	A	20040727	BR 2003-4218	20030329
				DE 2002-10214675	A 20020403
				WO 2003-EP3294	W 20030329
	EP 1492759	A2	20050105	EP 2003-732270	20030329
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				


OS CASREACT 139:307907; MARPAT 139:307907
GI



I



II



III

AB The invention relates to novel alicyclic esters I [R1 = Me; R2, R4 = H; R3 = H, Me; R5, R6 = H, Me; Y = CR7R8OC(:O)R9; R7, R8 = H, Me; R9 = C1-5-alkyl, C2-5-alkylene; or R1, R2 = Me, Et; R3, R4 = H, Me; R5R6 = O; Y = CR7R8OC(:O)R9; or R1, R2 = Me, Et; R4, R5, R6, R7 = H, Me; Y = CR7R8OC(:O)R9], methods for their production, for their use as odorous substances for perfumed products and for odorous substance mixts. containing the inventive compds. The procedure for the preparation of I is characterized by reaction of cyclohexylalkanols II with carboxylic acids [R9CO2CR7R8CO2H, R9CO2H or XCR7R8CO2H (X = OH, halogen)] anhydrides [(R9CO2)2O or (XCR7R8CO2)2O], or epoxides, III. Thus, I [R1 = Me, R2 - R4 = H, R6 = CHMe2, Y = O2CET] was prepared from 1-cyclohexylethanol via reaction with isobutylene oxide in cyclohexane containing BF3·OEt2, followed by reaction with (EtCO2)2O containing Et3N in the presence of catalytic DMAP. The odor of I [R1 = Me, R2 - R4 = H, R6 = CHMe2, Y = O2CET] was characterized (perceptible rose bloom note).

L8 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of cycloalkanecarboxylic acid derivatives for use as
fragrances with musk characteristics
AN 2002:925306 CAPLUS
DN 138:4714
TI Preparation of cycloalkanecarboxylic acid derivatives for use as
fragrances with musk characteristics
IN Kraft, Philip; Cadalbert, Riccardo
PA Givaudan SA, Switz.
SO Eur. Pat. Appl., 13 pp.
CODEN: EPXXDW
DT Patent


LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1262474	A1	20021204	EP 2001-113377	20010601
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	WO 2002096852	A1	20021205	WO 2002-CH282	20020530
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			EP 2001-113377	A 20010601
	EP 1392640	A1	20040303	EP 2002-727147	20020530
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
	BR 2002010098	A	20040413	EP 2001-113377	A 20010601
				WO 2002-CH282	W 20020530
	JP 2004535412	T2	20041125	BR 2002-10098	20020530
				EP 2001-113377	A 20010601
	US 2004234568	A1	20041125	WO 2002-CH282	W 20020530
				JP 2003-500032	20020530
				EP 2001-113377	A 20010601
				WO 2002-CH282	W 20020530
				US 2004-478626	20040614
				EP 2001-113377	A 20010601
				WO 2002-CH282	W 20020530

OS MARPAT 138:4714

GI



I

AB Odorant cycloalkanecarboxylic acid esters, such as I [R1, R2 = H, Me; X = (CH₂)_n; n = 1, 2, 3; n plus number of carbon atoms in R1 and R2 is less than 5], with musk odor were prepared for use as ingredients in fine-fragrances and functional perfumery compns. Thus, I (R1 = R2 = H, X = CH₂) was prepared by O-alkylation of α ,3,3-trimethylcyclohexylmethanol with isobutylene oxide using EtAlCl₂ in toluene, and subsequent esterification of the resulting alc. with cyclopropanecarboxylic acid using DMAP and DCC in CH₂Cl₂. Fragrance formulations of the prep'd cycloalkanecarboxylic acid esters were presented.

RE.CNT 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN

TI A method for the calculation of odor character from molecular structure

AN 2002:610108 CAPLUS

DN 139:3137
 TI A method for the calculation of **odor** character from molecular structure
 AU Turin, L.
 CS Department of Physiology, University College London, London, WC1E 6BT, UK
 SO Journal of Theoretical Biology (2002), 216(3), 367-385
 CODEN: JTBIAP; ISSN: 0022-5193
 PB Elsevier Science Ltd.
 DT Journal
 LA English
 AB The relationship between mol. structure and **odor** character is one of the most complex structure-activity problems in biol. Despite over a century of effort, it remains unsolved, and synthesis of new odorants still proceeds largely by trial and error. In previous work, I have argued that the reason for this failure lies in a mistaken assumption, namely that mol. shape dets. **odor** character. Instead, I have taken up and extended an old idea (Dyson, 1938) according to which vertebrate olfactory receptors detect odorants by their mol. vibrations. I propose that the detection mechanism is inelastic electron tunnelling. If this is correct, there should be a correlation between the tunnelling vibrational spectra of odorants and their **odor** character. Here, using semi-empirical quantum chemical methods and a simple calcn. method for tunnelling mode intensities, I calculate the spectra of structurally diverse odorants belonging to various **odor** categories. With few exceptions, the calculated spectra of bitter almonds, musks, ambers, woods, sandalwoods and violets strongly correlate with **odor** character.

RE.CNT 28 THERE ARE 28 CITED REFERENCES AVAILABLE FOR THIS RECORD
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
 TI Preparation of esters with musky **odor** and their use in perfumery
 AN 2000:175782 CAPLUS
 DN 132:194523
 TI Preparation of esters with musky **odor** and their use in perfumery
 IN Williams, Alvin S.
 PA Firmenich S. A., Switz.
 SO PCT Int. Appl., 24 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2000014051	A1	20000316	WO 1999-IB1469	19990825
	W: CA, CN, IL, IN, JP, US			CH 1998-1841	A 19980909
	RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE			CA 1999-2309449	19990825
		AA	20000316	CH 1998-1841	A 19980909
				WO 1999-IB1469	W 19990825
EP	1047660	A1	20001102	EP 1999-936930	19990825
EP	1047660	B1	20030129		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI			CH 1998-1841	A 19980909
				WO 1999-IB1469	W 19990825
JP	2002524438	T2	20020806	JP 2000-568811	19990825
				CH 1998-1841	A 19980909
				WO 1999-IB1469	W 19990825
AT	231834	E	20030215	AT 1999-936930	19990825
				CH 1998-1841	A 19980909
				WO 1999-IB1469	W 19990825
ES	2192067	T3	20030916	ES 1999-936930	19990825
				CH 1998-1841	A 19980909

US 6384269

B1 20020507

US 2000-553376

20000420

US 38659

E 20041123

WO 1999-IB1469

A1 19990825

US 2002-326341

20021223

CH 1998-1841

A 19980909

WO 1999-IB1469



A1 19990825

US 2000-553376

A5 20000420

OS MARPAT 132:194523

GI



AB Cyclohexanemethanol esters I [R1, R2, R3, R4, R6, R7 = H, Me, Et; R5 = alkyl, alkoxy, alkenyl, acyloxy; X = CR₆R₇], which have a musky odor, were prepared for use as perfume additives. Thus, ester II was prepared with 79.5% yield by esterification of (+)-[S-(R*,S*)]- $\alpha,3,3$ -trimethylcyclohexanemethanol with EtCO₂CH₂COCl. Perfuming compns. of the prepared compds. were presented.

RE.CNT 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L8 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2005 ACS on STN
TI Synthesis of ethylene glycol mixed esters of alcanoic and cyclohexanoic acids

AN 1968:77820 CAPLUS

DN 68:77820

TI Synthesis of ethylene glycol mixed esters of alcanoic and cyclohexanoic acids

AU Zeinalov, B. K.; Radzhalov, D. T.; Dzhafarova, A. A.; Muradyan, N. V.

SO Azerbaidzhanskii Khimicheskii Zhurnal (1967), (2), 35-7

CODEN: AZKZAU; ISSN: 0005-2531

DT Journal

LA Russian

GI For diagram(s), see printed CA Issue.

AB Acyloxyethyl esters (I and II) (tabulated) of cyclohexane- and 1,1-methylcyclohexanecarboxylic acids (sic) were prepared in 60-70% yield by heating a stirred mixture of 0.5 mole of the resp. alcanoic acid, 0.5 mole β -hydroxyethyl cyclohexanecarboxylate (II) or β -hydroxyethyl 1-methyl-1-cyclohexanecarboxylate (IV) 1% by weight of H₂SO₄, and 150 ml. dry C₆H₆ at 85-90° for 3-8 hrs., with a H₂O aspirator in operation, distilling C₆H₆ from the neutralized, washed solution, and vacuum distilling

the

residue. The products were oily liqs. with pleasant ester odors and had good solubilizing and plasticizing properties. Poly(vinyl chloride) plasticized with these esters was distinguished by high elasticity. III, b₁ 5 110-12°, d₂₀ 1.0668, n_{20D} 1.4680, and IV, b₁ 105-6°, d₂₀ 1.0436, n_{20D} 1.4665, were prepared by condensing the resp. acids with ClCH₂CH₂OH in the presence of 40% aqueous alkali. [TABLE OMITTED]

=> file reg
COST IN U.S. DOLLARS
FULL ESTIMATED COST

SINCE FILE ENTRY	TOTAL SESSION
30.78	196.82

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-5.11	-5.11

FILE 'REGISTRY' ENTERED AT 06:44:54 ON 19 JUL 2005
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2005 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file provided by InfoChem.

STRUCTURE FILE UPDATES: 17 JUL 2005 HIGHEST RN 855596-49-5
 DICTIONARY FILE UPDATES: 17 JUL 2005 HIGHEST RN 855596-49-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH JANUARY 18, 2005

Please note that search-term pricing does apply when conducting SmartSELECT searches.

 * The CA roles and document type information have been removed from *
 * the IDE default display format and the ED field has been added, *
 * effective March 20, 2005. A new display format, IDERL, is now *
 * available and contains the CA role and document type information. *

Structure search iteration limits have been increased. See HELP SLIMITS for details.

Experimental and calculated property data are now available. For more information enter HELP PROP at an arrow prompt in the file or refer to the file summary sheet on the web at:
<http://www.cas.org/ONLINE/DBSS/registryss.html>


```
=> e 1-Propanol, 2-(1-cyclohexyl-1-methylethoxy)-, propanoate/cn
E1      1      1-PROPANOL, 2-(1-BUTOXYETHOXY)-, (2R)-/CN
E2      1      1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY)-, ACETATE/CN
E3      1 --> 1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY)-, PROPAANOATE/CN
E4      1      1-PROPANOL, 2-(1-CYCLOHEXYLETHOXY)-2-METHYL-/CN
E5      1      1-PROPANOL, 2-(1-CYCLOHEXYLETHOXY)-2-METHYL-, ACETATE/CN
E6      1      1-PROPANOL, 2-(1-CYCLOHEXYLETHOXY)-2-METHYL-, PROPAANOATE/CN
E7      1      1-PROPANOL, 2-(1-CYCLOHEXYLPROPOXY)-2-METHYL-, ACETATE/CN
E8      1      1-PROPANOL, 2-(1-CYCLOHEXYLPROPOXY)-2-METHYL-, PROPAANOATE/CN
E9      1      1-PROPANOL, 2-(1-ETHOXYETHOXY)-/CN
E10     1      1-PROPANOL, 2-(1-ETHOXYETHOXY)-, (2R)-/CN
E11     1      1-PROPANOL, 2-(1-ETHOXYETHOXY)-, (2R)-(PARTIAL)-/CN
E12     1      1-PROPANOL, 2-(1-ETHOXYETHOXY)-, (2S)-/CN
```

```
=> e3
L9      1 "1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY)-, PROPAANOATE"/CN
```

=> d 19

```
L9      ANSWER 1 OF 1  REGISTRY  COPYRIGHT 2005 ACS on STN
RN      610769-92-1  REGISTRY
ED      Entered STN: 30 Oct 2003
CN      1-Propanol, 2-(1-cyclohexyl-1-methylethoxy)-, propanoate (9CI)
       (CA INDEX NAME)
FS      3D CONCORD
MF      C15 H28 O3
```

SR CA
LC STN Files: CA, CAPLUS, CASREACT




PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

=> search 19 exact full
L10 1 "1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY)-, PROPANOATE"/CN

=> d 110

L10 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2005 ACS on STN
RN 610769-92-1 REGISTRY
ED Entered STN: 30 Oct 2003
CN 1-Propanol, 2-(1-cyclohexyl-1-methylethoxy)-, propanoate (9CI)
(CA INDEX NAME)
FS 3D CONCORD
MF C15 H28 O3
SR CA
LC STN Files: CA, CAPLUS, CASREACT



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1907 TO DATE)
1 REFERENCES IN FILE CAPLUS (1907 TO DATE)

	SINCE FILE ENTRY	TOTAL SESSION
COST IN U.S. DOLLARS		
FULL ESTIMATED COST	13.31	210.13
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-5.11

FILE 'CAPLUS' ENTERED AT 06:45:43 ON 19 JUL 2005
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 19 Jul 2005 VOL 143 ISS 4
FILE LAST UPDATED: 18 Jul 2005 (20050718/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.


=> 110
L11 1 L10

=> d 111 ti fbib abs


L11 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN
TI Methods for the production of novel alicyclic esters having a musky smell
AN 2003:796643 CAPLUS
DN 139:307907
TI Methods for the production of novel alicyclic esters having a musky smell
IN Eh, Marcus
PA Symrise GmbH & Co. KG, Germany
SO PCT Int. Appl., 45 pp.
CODEN: PIXXD2
DT Patent
LA German
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2003082799	A2	20031009	WO 2003-EP3294	20030329
	WO 2003082799	A3	20031231		
	W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
	DE 10214675	A1	20031016	DE 2002-10214675	A 20020403
	BR 2003004218	A	20040727	BR 2003-4218	20030329
				DE 2002-10214675	A 20020403
				WO 2003-EP3294	W 20030329
	EP 1492759	A2	20050105	EP 2003-732270	20030329
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			DE 2002-10214675	A 20020403
				WO 2003-EP3294	W 20030329


OS CASREACT 139:307907; MARPAT 139:307907
GI



I



II



III

AB The invention relates to novel alicyclic esters I [R1 = Me; R2, R4 = H; R3 = H, Me; R5, R6 = H, Me; Y = CR7R8OC(:O)R9; R7, R8 = H, Me; R9 = C1-5-alkyl, C2-5-alkylene; or R1, R2 = Me, Et; R3, R4 = H, Me; R5R6 = O; Y = CR7R8OC(:O)R9; or R1, R2 = Me, Et; R4, R5, R6, R7 = H, Me; Y = CR7R8OC(:O)R9], methods for their production, for their use as odorous substances for perfumed products and for odorous substance mixts. containing the inventive compds. The procedure for the preparation of I is characterized by reaction of cyclohexylalkanols II with carboxylic acids [R9CO2CR7R8CO2H, R9CO2H or XCR7R8CO2H (X = OH, halogen)] anhydrides [(R9CO2)2O or (XCR7R8CO2)2O], or epoxides, III. Thus, I [R1 = Me, R2 - R4 = H, R6 = CHMe2, Y = O2CET] was prepared from 1-cyclohexylethanol via reaction with isobutylene oxide in cyclohexane containing BF3·OEt2, followed by reaction with (EtCO2)2O containing Et3N in the presence of catalytic DMAP. The odor of I [R1 = Me, R2 - R4 = H, R6 = CHMe2, Y = O2CET] was characterized (perceptible rose bloom note).

=> d his

(FILE 'HOME' ENTERED AT 06:25:36 ON 19 JUL 2005)

FILE 'REGISTRY' ENTERED AT 06:27:51 ON 19 JUL 2005

L1 STRUCTURE UPLOADED
 L2 50 SEARCH L1 SSS SAM
 L3 STRUCTURE UPLOADED
 L4 0 SEARCH L3 SSS SAM
 L5 74 SEARCH L3 SSS FULL
 SAVE TEMP L5 RAWFINDS/A

FILE 'CAPLUS' ENTERED AT 06:33:54 ON 19 JUL 2005

L6 21 L5
 SAVE TEMP ODORREFS/A L6
 L7 69532 ODOR
 L8 7 L6 AND L7

FILE 'REGISTRY' ENTERED AT 06:44:54 ON 19 JUL 2005
E 1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY)-, PROPAANOATE/CN

L9 1 E3
L10 1 SEARCH L9 EXACT FULL

FILE 'CAPLUS' ENTERED AT 06:45:43 ON 19 JUL 2005

L11 1 L10

=> 16 not 18

L12 14 L6 NOT L8

=> d 112 10-14 ti fbib abs

L12 ANSWER 10 OF 14 CAPLUS COPYRIGHT 2005 ACS on STN

TI Yellow dyes, ink compositions containing the same, and thermal transfer recording sheets using the same, giving high-density stable images resistant to heat, light, or humidity

AN 1998:693003 CAPLUS

DN 130:14995

TI Yellow dyes, ink compositions containing the same, and thermal transfer recording sheets using the same, giving high-density stable images resistant to heat, light, or humidity

IN Ogiso, Akira; Shimokawa, Yasushi; Ito, Naoto

PA Mitsui Chemicals Inc., Japan

SO Jpn. Kokai Tokkyo Koho, 17 pp.

CODEN: JKXXAF

DT Patent


LA Japanese

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 10287056	A2	19981027	JP 1997-98915	19970416
	JP 3582954	B2	20041027	JP 1997-98915	19970416

OS MARPAT 130:14995

GI



AB The title dyes have the general formula I, wherein R1, R2 = C1-10 (un)substituted alkyl, alkoxyalkyl; R3, R4 = H, Me; R5 = H, halogen, (un)substituted alkyl, alkoxy; n = 0, 1. An ink comprised I (R1 = R2 = Et; R3 = R5 = H; n = 0) 3, polybutyral resin 4.5, MEK 46.25, and toluene 46.25 parts.

L12 ANSWER 11 OF 14 CAPLUS COPYRIGHT 2005 ACS on STN

TI Branched bismethacrylates based on bis-GMA - a systematic route to low shrinkage composites

AN 1997:579234 CAPLUS

DN 127:221335

TI Branched bismethacrylates based on bis-GMA - a systematic route to low shrinkage composites

AU Holter, Dirk; Frey, Holger; Mulhaupt, Rolf
CS Institut fur Makromolekulare Chemie und Freiburger
Materialforschungszentrum (FMF) der Albert-Ludwigs-Universitat, Freiburg
i. Br., D-79104, Germany
SO Polymer Preprints (American Chemical Society, Division of Polymer
Chemistry) (1997), 38(2), 84-85
CODEN: ACPPAY; ISSN: 0032-3934
PB American Chemical Society, Division of Polymer Chemistry
DT Journal
LA English
AB Bismethacrylate monomers were prepared by esterification of bisphenol A
diglycidyl methacrylate (Bis-GMA) with different aliphatic and aromatic acids.
The relationship between mol. structure, viscosity, volume shrinkage and
mech. properties was studied. Variation of the side groups attached to
the Bis-GMA scaffold allows to decrease viscosity and volume shrinkage
concurrently. To obtain reasonable mech. properties Ph rings were
introduced into the side groups. 4-Phenylbutyric acid derivatized Bis-GMA
shows a viscosity of 51 Pas. The resp. composite with 50 vol% of a glass
filler reveals a volume shrinkage of 1.5% and an E-modulus of 6,300 MPa.

L12 ANSWER 12 OF 14 CAPLUS COPYRIGHT 2005 ACS on STN
TI Lipase-catalyzed resolution of 3-(aryloxy)-1,2-propanediol derivatives.
Towards an improved active site model of *Pseudomonas cepacia* lipase (Amano
PS)
AN 1995:663901 CAPLUS
DN 123:339233
TI Lipase-catalyzed resolution of 3-(aryloxy)-1,2-propanediol derivatives.
Towards an improved active site model of *Pseudomonas cepacia* lipase (Amano
PS)
AU Theil, Fritz; Lemke, Karin; Ballschuh, Sibylle; Kunath, Annamarie; Schick,
Hans
CS Inst. Angewandte Chemie Berlin-Adlershof e.V., Berlin, D-12484, Germany
SO Tetrahedron: Asymmetry (1995), 6(6), 1323-44
CODEN: TASYE3; ISSN: 0957-4166
PB Elsevier
DT Journal
LA English
OS CASREACT 123:339233
AB A variety of 3-(aryloxy)-1,2-propanediol derivs. with different
substituents on the aromatic ring or at the primary hydroxy group were used
as substrates for kinetic resolution by transesterification with vinyl
acetate catalyzed by lipase from *Pseudomonas cepacia* (Amano PS). Derivs.
with substituents in the para position of the aromatic ring were accepted as
substrates and resolved with high enantioselectivity. The corresponding
derivs. with substituents in the ortho position were much worse substrates
for lipase PS or even non-substrates if the substituent was sufficiently
space-filling, e.g., tert-Bu, Ph, benzyl or benzoyl. If the primary
hydroxy group was substituted by unbranched long-chain acyl residues, very
good substrates resulted. In contrast, derivs. with sterically crowded
residues at the primary hydroxy group (e.g., pivaloyl,
tert-butyldimethylsilyl, methanesulfonyl, p-toluenesulfonyl, or trityl)
were non-substrates for lipase PS.

L12 ANSWER 13 OF 14 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of 4-cycloalkyl-3-oxapentyl alkanoates as perfume fragrances
AN 1992:407513 CAPLUS
DN 117:7513
TI Preparation of 4-cycloalkyl-3-oxapentyl alkanoates as perfume fragrances
IN Giersch, Wolfgang Klaus; Schulte-Elte, Karl Heinrich
PA Firmenich S. A., Switz.
SO Eur. Pat. Appl., 15 pp.
CODEN: EPXXDW
DT Patent
LA French
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 472966	A1	19920304	EP 1991-113240	19910807
	EP 472966	B1	19940928		
	R: CH, DE, FR, GB, LI, NL				
	US 5166412	A	19921124	CH 1990-2799	A 19900828
				US 1991-741027	19910806
	JP 06072952	A2	19940315	CH 1990-2799	A 19900828
	JP 2974834	B2	19991110	JP 1991-214881	19910827
				CH 1990-2799	A 19900828
OS	MARPAT 117:7513				
AB	RCHMeOCR1R2CR3R4O2CR5 (R = 3,3-dimethylcyclopentyl, -cyclohexyl; when R1 = R2 = H, R3 and/or R4 = Me; when R3 = R4 = H, R1 and/or R2 = Me; R5 = alkyl) were prepared. Thus, 1-(3,3-dimethyl-1-cyclohexyl)-1-ethanone was ketalized by HOCH2CHMeOH and the dioxolane product reduced with Dibal to give RCHMeOCHMeCH2OR4 (I; R = 3,3-dimethylcyclohexyl) (II; R4 = H) and RCHMeOCH2CHMeOR4 (III; R same as I) (IV; R4 = H) as a mixture which was treated with EtCOCl to give II and IV (R4 = EtCO in each) as mixts. of diastereomers. Perfume formulations comprising title compds. are given.				
L12	ANSWER 14 OF 14 CAPLUS COPYRIGHT 2005 ACS on STN				
TI	Photopolymerizing compositions				
AN	1983:603607 CAPLUS				
DN	99:203607				
TI	Photopolymerizing compositions				
PA	Mitsubishi Chemical Industries Co., Ltd., Japan				
SO	Jpn. Kokai Tokkyo Koho, 7 pp.				
	CODEN: JKXXAF				
DT	Patent				
LA	Japanese				
FAN.CNT 1					
	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	JP 58134629	A2	19830810	JP 1982-16761	19820204
				JP 1982-16761	19820204
GI					



AB Photopolymg. compns. contain (1) a binder polymer having monomer unit of the general formula $\text{CH}_2\text{CRCO}_2\text{C}_2\text{H}_4\text{OCOZCO}_2\text{H}$ (I) [R = H, Me; Z = II, III, IV, $(\text{CH}_2)_n$, CH:CR2 (R1 = H, CO2H; R2 = H, Me; n = 2, 3)], (2) an unsatd. compound polymerizable by addition having ≥ 1 ethylenically unsatd. bonds, and (3) a photopolymn. initiator. The composition provides photosensitive materials with good adhesivity and abrasion resistance. Thus, Me methacrylate 85.5, isophthalic acid mono- β -methacryloxyethyl ester 12.5 and benzoyl peroxide 0.5 were reacted and purified to obtain a copolymer suitable as a binder polymer. An Al support was coated first with a mixture of the binder polymer, pentaerythritol triacrylate, and a photopolymn. initiator and subsequently overcoated with a poly(vinyl alc.) layer to give a photopolymn. imaging material. The material showed excellent performance.

=> logoff hold
COST IN U.S. DOLLARS

SINCE FILE TOTAL
ENTRY SESSION

FULL ESTIMATED COST	18.15	228.28
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-4.38	-9.49

SESSION WILL BE HELD FOR 60 MINUTES
 STN INTERNATIONAL SESSION SUSPENDED AT 06:48:34 ON 19 JUL 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623PAZ

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
 SESSION RESUMED IN FILE 'CAPLUS' AT 07:14:36 ON 19 JUL 2005
 FILE 'CAPLUS' ENTERED AT 07:14:36 ON 19 JUL 2005
 COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	18.15	228.28
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-4.38	-9.49

=> d his

(FILE 'HOME' ENTERED AT 06:25:36 ON 19 JUL 2005)

FILE 'REGISTRY' ENTERED AT 06:27:51 ON 19 JUL 2005

L1 STRUCTURE UPLOADED
 L2 50 SEARCH L1 SSS SAM
 L3 STRUCTURE UPLOADED
 L4 0 SEARCH L3 SSS SAM
 L5 74 SEARCH L3 SSS FULL
 SAVE TEMP L5 RAWFINDS/A

FILE 'CAPLUS' ENTERED AT 06:33:54 ON 19 JUL 2005

L6 21 L5
 SAVE TEMP ODORREFS/A L6
 L7 69532 ODOR
 L8 7 L6 AND L7

FILE 'REGISTRY' ENTERED AT 06:44:54 ON 19 JUL 2005
 E 1-PROPANOL, 2-(1-CYCLOHEXYL-1-METHYLETHOXY) -, PROPANOATE/CN

L9 1 E3
 L10 1 SEARCH L9 EXACT FULL

FILE 'CAPLUS' ENTERED AT 06:45:43 ON 19 JUL 2005

L11 1 L10
 L12 14 L6 NOT L8

=> fragr?
 L13 13878 FRAGR?

=> l6 and l13
 L14 3 L6 AND L13

=> l14 not 18

L15 1 L14 NOT L8

=> d 115 ti fbib abs

L15 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2005 ACS on STN
TI Preparation of 4-cycloalkyl-3-oxapentyl alkanoates as perfume
fragrances
AN 1992:407513 CAPLUS
DN 117:7513
TI Preparation of 4-cycloalkyl-3-oxapentyl alkanoates as perfume
fragrances
IN Giersch, Wolfgang Klaus; Schulte-Elte, Karl Heinrich
PA Firmenich S. A., Switz.
SO Eur. Pat. Appl., 15 pp.
CODEN: EPXXDW
DT Patent
LA French
FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 472966	A1	19920304	EP 1991-113240	19910807
	EP 472966	B1	19940928		
	R: CH, DE, FR, GB, LI, NL			CH 1990-2799	A 19900828
	US 5166412	A	19921124	US 1991-741027	19910806
				CH 1990-2799	A 19900828
	JP 06072952	A2	19940315	JP 1991-214881	19910827
	JP 2974834	B2	19991110	CH 1990-2799	A 19900828

OS MARPAT 117:7513
AB RCHMeOCR1R2CR3R4O2CR5 (R = 3,3-dimethylcyclopentyl, -cyclohexyl; when R1 = R2 = H, R3 and/or R4 = Me; when R3 = R4 = H, R1 and/or R2 = Me; R5 = alkyl) were prepared. Thus, 1-(3,3-dimethyl-1-cyclohexyl)-1-ethanone was ketalized by HOCH2CHMeOH and the dioxolane product reduced with Dibal to give RCHMeOCHMeCH2OR4 (I; R = 3,3-dimethylcyclohexyl) (II; R4 = H) and RCHMeOCH2CHMeOR4 (III; R same as I) (IV; R4 = H) as a mixture which was treated with EtCOCl to give II and IV (R4 = EtCO in each) as mixts. of diastereomers. Perfume formulations comprising title compds. are given.

=> d cost

COST IN U.S. DOLLARS

SINCE FILE ENTRY	TOTAL SESSION
---------------------	------------------

CONNECT CHARGES

4.68

16.37

NETWORK CHARGES

0.72

2.70

SEARCH CHARGES

1.89

173.88

DISPLAY CHARGES

18.55

43.02

FULL ESTIMATED COST

25.84

235.97

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE ENTRY	TOTAL SESSION
---------------------	------------------

CA SUBSCRIBER PRICE

-5.11

-10.22

IN FILE 'CAPLUS' AT 07:18:43 ON 19 JUL 2005

=> logoff hold

COST IN U.S. DOLLARS

SINCE FILE ENTRY	TOTAL SESSION
---------------------	------------------

FULL ESTIMATED COST

25.84

235.97

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE ENTRY	TOTAL SESSION
---------------------	------------------

CA SUBSCRIBER PRICE

-5.11 -10.22

SESSION WILL BE HELD FOR 60 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 07:19:00 ON 19 JUL 2005

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1623PAZ

PASSWORD:

* * * * * RECONNECTED TO STN INTERNATIONAL * * * * *
SESSION RESUMED IN FILE 'CAPLUS' AT 08:01:54 ON 19 JUL 2005
FILE 'CAPLUS' ENTERED AT 08:01:54 ON 19 JUL 2005
COPYRIGHT (C) 2005 AMERICAN CHEMICAL SOCIETY (ACS)

COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	25.84	235.97
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-5.11	-10.22
=> logoff hold		
COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	25.84	235.97
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-5.11	-10.22

SESSION WILL BE HELD FOR 60 MINUTES
STN INTERNATIONAL SESSION SUSPENDED AT 08:02:06 ON 19 JUL 2005